

SCOTLAND HEALTH CARE SYSTEM
Clinical Privileges Delineation Form
Specialty: Obstetrics/Gynecology

Physician Name: _____

I. **General Privileges**

By virtue of being granted any level of clinical privileges, a physician is automatically granted the following privileges:

Performing physical examinations and ordering and interpreting of diagnostic studies that are normally considered part of the practice of obstetrics/gynecology, including, but not limited to, laboratory, diagnostic imaging, and electrocardiographic studies.

Prescribing and administering medications normally considered part of the practice of obstetrics/gynecology.

Requesting consultations and technical procedures to be performed by other physicians and qualified consultants/technicians.

At the time of a clinical emergency, the provision of whatever care the physician believes to be indicated to prevent loss of life or serious injury to a patient.

II. **Specific Privileges: Clinical**

Physicians granted clinical privileges in obstetrics/gynecology must demonstrate knowledge and skills usually achieved only through training sufficient to attain eligibility for Board certification in obstetrics/gynecology, or equivalent experience. (NOTE: if sufficient training is not documented, specific experience must be outlined in writing).

OB/GYN

| Procedure | Requested | Procedures | Requested |
|--|-----------|--|-----------|
| D&C - Diagn./Therapeutic | | Vaginal Suspension Operations | |
| I&D - Bartholin duct absc. | | Salpingostomy | |
| Bartholin duct cystectomy | | Umbilical hernia repair | |
| Marsupialization - Bartholin Cyst | | Urethral diverticulectomy | |
| Biopsy of vulva | | Excision periurethral absc. | |
| Biopsy of cervix | | Repair urethro-vaginal fist. | |
| Biopsy of vagina | | LaForte vaginal repair | |
| Conization of cervix - cold knife | | Repair/recto-vaginal fistula | |
| LEEP procedure | | Repair/vesico-vaginal fist. | |
| Perineorrhaphy | | Sturmdorf repair of cervix | |
| Perineotomy | | Meckel's diverticulum + Appendectomy | |
| Repair - rectocele | | Wedge resect. - ovaries | |
| Repair - enterocoele | | Hydatid mole evacuat. | |
| Repair - cysto urethrocoele | | Salpingoplasty | |
| Excsn. - Skenes duct cyst | | Tubal implant - uterus | |
| Excsn./ fulguration of urethral caruncle | | Tubal anastomosis | |
| Vulvectomy - simple | | Closure - vaginal fistula | |
| Hysterectomy - vaginal | | Evacuation of pelvic absc. | |
| Hysterectomy - laparoscopic | | Evisceration repair | |
| Hysterectomy, abdominal, Total w/wo adnexae | | Removal - foreign body vagina & uterus | |
| Subtotal w/wo adnexae | | Perineoplasty | |
| Uterine Suspension | | Burch urethropexy | |
| Pre-sacral neurectomy | | Laparoscopic Burch urethropexy | |
| Marshall-Marchetti pubo-vesico-urthral suspen. | | Needle urethropexy | |
| Hysterectomy, radical, Wertheim | | Laparoscopy - Diagnostic | |
| Colpectomy | | - Operative | |
| Pessary insertion | | Sterilizations - Abdominal | |
| Colpotomy-exploratory | | - Laparoscopic | |
| Salpingectomy | | Myomectomy | |

| Procedure | Requested | Procedure | Requested |
|---|-----------|--|-----------|
| Oophorectomy | | Metroplasty | |
| Hypogastric Artery Ligation | | Hysteroscopy-Diagnostic | |
| Hymenotomy | | - Operative | |
| Hymenectomy | | Hysterotomy | |
| Cerclage | | Perineorrhaphy | |
| Repair surgical incision: bladder, bowel | | Culdocentesis | |
| Pelvic & Para Aortic lymphadenectomy | | Trachelectomy | |
| Ureteral repair | | Incisional hernia repair | |
| Ureteral transplant emergency | | Hypogastric Artery Ligation | |
| Laser - Intraperitoneal | | Ectopic Pregnancy- salping- ectomy, salpingostomy | |
| -Extraperitoneal Lower genital track | | Treatment of medical com- plications of obstetrics - heart, lungs, kidney, anemia, diabetes, etc. | |
| Endometrial biopsy | | Repair of 3rd & 4th degree lacerations | |
| Gynecologic U/S | | | |
| OBSTETRICS | | Cudocentensis | |
| Spont. delivery, vertex presentation | | Piper forceps application to after coming head | |
| * emergency | | Repair uterine lacerations | |
| * non-emergency | | Repair-perineal lacerations | |
| Episiotomy & repair | | Abdominal pregnancy | |
| Breech delivery: * Multipara | | Colpotomy | |
| * Primigravida | | Amniotomy | |
| * Frank | | Anesthesia-pudanal, block, local | |
| * Footling | | *General - emergency | |
| Multiple Pregnancy | | * Spinal - elective | |
| Outlet Forceps | | * Epidural - elective | |
| Low Forceps: - nonrotational | | Evacuation of vulvar hematomata | |
| - rotational | | Excsn. - vulvar lesions at delivery | |
| Mid forceps delivery w/wo rotation | | Excision-vaginal cysts | |

| Procedure | Requested | Procedure | Requested |
|---|-----------|--|-----------|
| Vacuum Extraction | | Uterine packing | |
| Pre-eclampsia - mild | | Curettage uterine | |
| - severe | | Suction cutterage | |
| Eclampsia | | Abortion - 1st Tri.mester | |
| Cesarean Section: | | Abortion - 2nd Trimester | |
| -Classical | | Post-partum Care | |
| -Low cervical | | Resuscitation of newborn | |
| -Extrperitoneal | | Circumcision of infant | |
| External cephalic version-singleton | | Cervical biopsy during pregnancy- conization of cervix | |
| Version & extraction of twins | | Repair of incompetent or cervical os circlage | |
| Manual removal of placenta | | Hemorrhoid excision / incision | |
| Repair - cervical lacerations | | Inversion of uterus | |
| Cesarean hysterectomy | | Amniocentesis | |
| Uterine artery ligation | | Application of leads for internal monitoring of fetus | |
| Mgmt. of fetal death in utero-intra- amniotic injection of hypertonic solutions | | Application - transducers for internal monitoring of fetus | |
| Mgmt. of Prostaglandin gel insertion in utero-intra- amniotic injection of hypertonic solutions | | Mgmt. of Oxytocin for augmentation or induction - labor | |
| Induction of labor: * Medical | | Mgmt. of Prostaglandin gel for augmentation or induction - labor | |
| Surgical* | | Manage high risk obstetrical patients in labor | |
| Missed Abortion - 1st Trimester | | Manage post partum hemorrhage | |
| Missed Abortion - 2nd Trimester | | Manage choriocarcinoma | |
| Management of: * Premature rupture of membranes | | Manage sepsis during pregnancy | |
| *Premature labor | | Vaginal Laceration Repair | |
| * Antepartum fetal death | | Interpretation NST, CST, NCST, OCT, BPP | |
| * Amniotic fluid - pulmonary embolus | | Obstetric ultrasound | |
| Application of leads for internal monitoring of fetus | | | |
| Application - transducers for internal monitoring of fetus | | | |

I hereby request the clinical privileges listed above.

Applicant Signature: _____ Date: _____