No aspirin or salicylic acid in children of any age Date: Age: Weight:

- Keep all medicines out of reach of children
- No Ibuprofen for children less than 6 months old
- Ibuprofen, Children's Motrin® and Children's Advil® are all the same
- Give the dose for your child's weight as listed below
- Give every 6 hours for fever more than 100.4 degrees

Weight (pounds)	Infant Drops 50 mg per 1.25 ml	Children's 100 mg per 5 ml	Junior 100 mg per tablet
12 to 17	if your child is over 6 months old, call doctor for dose		
18 to 23	1.875 ml		
24 to 35		5 ml (1 tsp)	1 tablet
36 to 47		7.5 ml	1½ tablets
48 to 59		10 ml (2 tsp)	2 tablets
60 to 71		12.5 ml	2½ tablets
72 to 95		15 ml (3 tsp)	3 tablets
more than 95		20 ml (4 tsp)	4 tablets

Give your child the right dose

- Be sure to read the right dose for your child
- Only use the device you get with the medicine to give the right dose

tsp = teaspoon

ml = milliliter

1 tsp = 5 ml



Dosing information for Acetaminophen and Ibuprofen from Tylenol for Health Professionals https://www.tylenolprofessional.com/dosage.html

Approved by Atrium Health Antibiotic Stewardship Committee, Fall 2018

Atrium Health complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability or sex.

ATENCIÓN: si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística. Llame al 1-800-821-1535.

CHÚ Ý: Nêu ban nói Tiêng Việt, có các dịch vu hô tro ngôn ngû miên phí dành cho ban. Goi sô 1-800-821-1535.

