

## Atrium Health Infusion Center

Referral Status:	☐ New Start ☐ Order Change ☐ Renewal		
Preferred Location:	Atrium Health Infusion Center Concord Fax: 704-468-3401		
	Atrium Health Infusion Center Pineville Fax: 704-468-3401		
	Atrium Health Infusion Center Southpark Fax: 704-468-3401		
	Atrium Health Infusion Center Huntersville Fax: 704-468-3401		
	Atrium Health Infusion Center Kenilworth, a facility of CMC Fax: 704-512-5390		
	Atrium Health Infusion Center Abbey Place, a facility of CMC Fax: 704-512-5390		
	Atrium Health Infusion Center Cabarrus, a facility of CMC Fax: 704-512-5390		

Vistide® (cidofivir) Infusion Order (Revised 11/7/2025)

All orders with a $$ will be placed.					
Patient Demographics:					
Patient Name:	Date of Birth:	MRN:			
Address:		'			
City:	State:	Zip Code:			
Allergies: (please list all allergies or attach list)	!	, .			
□ NKDA					
Diagnosis: (Complete the 2nd and/or 3rd Dig	its of the ICD-10)				
☐ ICD-10:	· · · · · · · · · · · · · · · · · · ·	Other:			
Required Documentation: (required prior to s	cheduling )				
Patient Demographic Sheet	0,	If the patient is new to the ordered therapy, indicate washout from previous therapy:			
Copy of Insurance Card (front and back)		, and parameter than to ano oracle and apply, and one and apply			
Most Recent Labs (must include labs pertinent to medication ordered )		☐ No Washout Needed			
Consult Note or last 2 Office Visits with referring provider or APP		If the patient is currently on the therapy, indicate date of last infusion:			
Complete Medication List -		Next infusion due date:			
Include all tried and failed meds					
		If this is an order change only, indicate if the current therapy should be administered until			
		insurance approval is received for the new request.			
Diagnostic Studies Portinent to Medication Or	lorod				
Diagnostic Studies Pertinent to Medication Ord	iereu	☐ Yes	□ No		
Nursing Communication:					
✓ Start PIV/Access CVC and flush device per approved Atrium Health protocol.					
Obtain vital signs PRE-treatment and POST-					
Monitor patient for signs of reaction for 30n			s PRN IF previous signs of reaction observed.		
Pre-Medications:		·			
☐ Acetaminophen (Tylenol) 650mg PO ONCE					
Diphenhydramine (Benadryl) 25mg PO ONCE					
Diphenhydramine (Benadryl) 25mg IV ONCE					
☐ Loratadine (Claritin) 10mg PO ONCE					
Methylprednisolone Sodium Succinate (Solu-Medrol) 125mg IV ONCE					
Hydration:					
☐ Pre-hydration: Sodium Chloride 0.9% 1000mL IV over 60 minutes every 2 weeks					
Post-hydration: Sodium Chloride 0.9% 1000mL IV over 60 minutes every 2 weeks					
Infusion Therapy:					
Cidofivir (Vistide) 0.5mg/kg IV at 200mL/hr (over 60 minutes) every 2 weeks					
Supportive Care Medications:					
Acetaminophen (Tylenol) 650mg PO ONCE PRN mild pain (1-3) or moderate pain (4-6). Give first if not given as a pre-medication.					
✓ Ibuprofen (Motrin) 800mg PO ONCE PRN mild pain (1-3) or moderate pain (4-6). Give second after acetaminophen.					
Ondansetron (Zofran) 4mg IV ONCE PRN nausea/vomiting.					
Hypersensitivity Protocol:					
Initiate Atrium Health approved hypersensitivity protocol in the event of an acute adverse or anaphylactic infusion/injection reaction. The hypersensitivity					
protocol can be found on the Atrium Health Infusion Center website at atriumhealth.org/infusion.					
Prescriber Information:					
Provider Name:		Phone:	Fax:		
Practice Name:		NPI:			
Address:		Office Contact:			
		Office Contact Phone Number:			
Physician Signature: (Order expires 12 months from date of signature ) No Stamp Signatures Accepted					
Signature:		Date:			
orginaturo.		54.0.			