

Atrium Health Infusion Center

Referral Status:	New Start ☐ Order Change ☐ Renewal
Preferred Location:	Atrium Health Infusion Center Concord Fax: 704-468-3401
	Atrium Health Infusion Center Pineville Fax: 704-468-3401
	Atrium Health Infusion Center Southpark Fax: 704-468-3401
	Atrium Health Infusion Center Huntersville Fax: 704-468-3401
	☐ Atrium Health Infusion Center Kenilworth, a facility of CMC Fax: 704-512-5390
	☐ Atrium Health Infusion Center Abbey Place, a facility of CMC Fax: 704-512-5390
	Atrium Health Infusion Center Cabarrus, a facility of CMC Fax: 704-512-5390

Cinqair® (reslizumab) Infusion Order (Revised 11/5/2025)

All orders with a $$ will be placed.			
Patient Demographics:			
Patient Name:	Date of Birth:	MRN:	
Address:	·	•	
City:	State:	Zip Code:	
Allergies: (please list all allergies or	attach list)	•	
■ NKDA			
Diagnosis: (Complete the 2nd and/	or 3rd Digits of the ICD-10)		
J45.50 - Severe persistent asthma, uncomplicated		☐ J45.51 - Severe persistent asthma with (acute) exacerbation	
J45.52 - Severe persistent asthma with status asthmaticus		Other:	
Required Documentation: (require	d prior to scheduling)		
Patient Demographic Sheet		If the patient is new to the ordered therapy, indicate washout from previous therapy:	
Copy of Insurance Card (front and back)			
Most Recent Labs (must include labs pertinent to medication ordered)		No Washout Needed	
		If the patient is currently on the therapy, indicate date of last infusion:	
Consult Note or last 2 Office Visits with referring provider or APP		Next infusion due date:	
Complete Medication List -			
Include all tried and failed meds		If this is an order change only, indicate if the current therapy should be administered until	
Diagnostic Studies Pertinent to Med	ication Ordered	in this is all order change only, indicate if the current therapy should be administered until	
Diagnostic Studies Fertillelit to Med	ication Ordered	☐ Yes ☐ No	
Nursing Communication:			
	device per approved Atrium Health (protocol	
Start PIV/Access CVC and flush device per approved Atrium Health protocol. Obtain vital signs DPE treatment, halfway through treatment, and POST treatment.			
Obtain vital signs PRE-treatment, halfway through treatment, and POST-treatment. Monitor nation for signs of reaction for 20mins AFTER completion of first 3 treatments and subsequent treatments PRN if previous signs of reaction observed.			
Monitor patient for signs of reaction for 30mins AFTER completion of first 3 treatments and subsequent treatments PRN if previous signs of reaction observed.			
Pre-Medications: (Administer all pre-medications 30mins prior to treatment)			
Acetaminophen (Tylenol) 650mg PO ONCE			
Diphenhydramine (Benadryl) 25mg PO ONCE			
Diphenhydramine (Benadryl) 25mg IV ONCE			
Loratadine (Claritin) 10mg PO ONCE			
Methylprednisolone Sodium Succinate (Solu-Medrol) 125mg IV ONCE			
Infusion Therapy:			
Cinqair (reslizumab) 3mg/kg IV every 4 weeks			
- Administer over 50 minutes for first 3 infusions only, and then increase rate as tolerated to a minimum infusion rate of 20 minutes.			
Supportive Care Medications:			
Ondansetron (Zofran) 4mg IV ON	ICE PRN nasusea/vomiting.		
Hypersensitivity Protocol:			
		t of an acute adverse or anaphylactic infusion/injection reaction. The hypersensitivity protocol can	
be found off the Athum Health in	fusion Center website at atriumheal	lth.org/infusion.	
Prescriber Information:			
Provider Name:		Phone: Fax:	
Practice Name:		NPI:	
Address:		Office Contact:	
City, State, Zip: Office Contact Phone Number:			
Physician Signature: (Order expires 12 months from date of signature) No Stamp Signatures Accepted			
Signature:		Date:	