

Atrium Health Infusion Center

Referral Status:	☐ New Start ☐ Order Change ☐ Renewal
Preferred Location:	Atrium Health Infusion Center Concord Fax: 704-468-3401
	Atrium Health Infusion Center Pineville Fax: 704-468-3401
	Atrium Health Infusion Center Southpark Fax: 704-468-3401
	Atrium Health Infusion Center Huntersville Fax: 704-468-3401
	Atrium Health Infusion Center Kenilworth, a facility of CMC Fax: 704-512-5390
	Atrium Health Infusion Center Abbey Place, a facility of CMC Fax: 704-512-5390
	Atrium Health Infusion Center Cabarrus, a facility of CMC Fax: 704-512-5390

Nulojix® (belatacept) Infusion Order (Revised 11/7/2025)

All orders with a √ will be placed.					
Patient	t Demographics:				
Patient	Name:	Date of Birth:	MRN:		
Addres	ss:	·	·		
City:		State:	Zip Code:		
Allergie	es: (please list all allergie	es or attach list)	·		
□ NKDA					
Diagnosis: (Complete the 2nd and/or 3rd Digits of the ICD-10)					
Z94.0 - Kidney Transplant Status Other:					
Required Documentation: (required prior to scheduling)					
	Demographic Sheet		If the patient is new to the ordered therapy, indicate washout from previous therapy:		
Copy of Insurance Card (front and back)		nd back)			
Most Recent Labs (must include labs pertinent to medication ordered)		·	☐ No Washout Needed		
Consult Note or last 2 Office Visits with referring provider or APP		sits with referring provider or APP	If the patient is currently on the therapy, indicate date of last infusion:		
Complete Medication List -			Next infusion due date:		
Include	e all tried and failed meds	S	If this is an order change only, indicate if the current therapy should be administered until		
			insurance approval is received for the new request.		
Diagno	stic Studies Pertinent to	Medication Ordered	_ `		
			Yes No		
Nursin	g Communication:				
_		ush device per approved Atrium Health p			
		nent and POST-treatment. Obtain vital si			
		eaction for 30mins AFTER completion of	1st infusion and subsequent infusions PRN IF previous signs of reaction observed.		
	edications:				
	taminophen (Tylenol) 65				
	henhydramine (Benadryl)				
	henhydramine (Benadryl)				
Loratadine (Claritin) 10mg PO ONCE					
Methylprednisolone Sodium Succinate (Solu-Medrol) 125mg IV ONCE					
Infusion Therapy:					
Required for Weight-Based Dosing: Transplant Date: Patient Transplant Weight: - Dose is calculated on transplant weight unless weight varies by >10%, after which patient will be dosed on actual body weight					
Check here if dose is to be calculated with actual body weight					
		ulojix) 10mg/kg IV over 30 minutes			
	Day 1:	Dose:			
	Day 15:	Dose:			
	Day 29:	Dose:			
\vdash	Day 42:	Dose:			
<u> </u>	Day 57:	Dose:			
	ntenance Phase: Belata rtive Care Medications:	cept (Nulojix) 5mg/kg IV over 30 minutes	every 4 weeks		
			adorate pain (A.S.). Cive first if not given as a pre-modication		
			oderate pain (4-6). Give first if not given as a pre-medication.		
Ubuprofen (Motrin) 800mg PO ONCE PRN mild pain (1-3) or moderate pain (4-6). Give second after acetaminophen.					
Ondansetron (Zofran) 4mg IV ONCE PRN nausea/vomiting. Hypersensitivity Protocol:					
	-	yed bypereepoitivity protectlin the event	of an aguta advarga or anaphylactic influsion/injection reaction. The hyperconsitivity		
Initiate Atrium Health approved hypersensitivity protocol in the event of an acute adverse or anaphylactic infusion/injection reaction. The hypersensitivity protocol can be found on the Atrium Health Infusion Center website at atriumhealth.org/infusion.					
Prescriber Information:					
	er Name:		Phone: Fax:		
	e Name:		NPI:		
Addres			Office Contact:		
-	tate, Zip:		Office Contact Phone Number:		
Physician Signature: (Order expires 12 months from date of signature) No Stamp Signatures Accepted					
Signatu	ure:		Date:		