

Atrium Health Infusion Center

| Referral Status: | ☐ New Start ☐ Order Change ☐ Renewal | | |
|---------------------|---|--|--|
| Preferred Location: | ☐ Atrium Health Infusion Center Concord Fax: 704-468-3401 ☐ Atrium Health Infusion Center Pineville Fax: 704-468-3401 ☐ Atrium Health Infusion Center Southpark Fax: 704-468-3401 | | |
| | | | |
| | | | |
| | Atrium Health Infusion Center Huntersville Fax: 704-468-3401 | | |
| | Atrium Health Infusion Center Kenilworth, a facility of CMC Fax: 704-512-5390 | | |
| | Atrium Health Infusion Center Abbey Place, a facility of CMC Fax: 704-512-5390 | | |
| | Atrium Health Infusion Center Cabarrus, a facility of CMC Fax: 704-512-5390 | | |

| Spevigo® (spesolimab-sbzo) Infusion Order (Revised 11/7/2025) All orders with a $$ will be placed. | | | | |
|---|--|--|--|--|
| Patient Demographics: | | | | |
| Patient Name: | Date of Birth: | MRN: | | |
| Address: | Date of Birth. | 11111 | | |
| City: | State: | Zip Code: | | |
| Allergies: (please list all allergies or attach list) | State. | Zip Code. | | |
| | | | | |
| ☐ NKDA Diagnosis: (Complete the 2nd and/or 3rd Digits of the ICD-10) | | | | |
| L40.1 - Generalized Pustular Psoriasis | of the ICD-10) | | | |
| | | Other: | | |
| Required Documentation: (required prior to sch | reduting) | | | |
| Patient Demographic Sheet | | If the patient is new to the ordered therapy, indicate washout from previous therapy: | | |
| Copy of Insurance Card (front and back) | | | | |
| Most Recent Labs (must include labs pertinent to medication ordered) | | ☐ No Washout Needed | | |
| Consult Note or last 2 Office Visits with referring provider or APP | | If the patient is currently on the therapy, indicate date of last infusion: Next infusion due date: | | |
| Complete Medication List - | | | | |
| Include all tried and failed meds | | | | |
| | | If this is an order change only, indicate if the current therapy should be administered until | | |
| | | insurance approval is received for the new request. | | |
| Diagnostic Studies Pertinent to Medication Order | red | ☐ Yes ☐ No | | |
| Treatment Parameters | | | | |
| | | | | |
| Hold treatment and notify provider IF: | | | | |
| - Temperature is GREATER THAN 100oF; | | | | |
| | - Patient complains of symptoms of acute viral or bacterial infection; | | | |
| - Patient is taking an antibiotic for current infection. | | | | |
| Required lab results: PPD/Quatiferon Gold PRIOR to first treatment. (fax labs with order) | | | | |
| | | | | |
| - Hold Tx and Notify Provider IF: | | | | |
| - PPD/Quantiferon Gold: POSITIVE result or not on file. | | | | |
| Nursing Communication: | | | | |
| ✓ Start PIV/Access CVC and flush device per approved Atrium Health protocol. | | | | |
| Obtain vital signs every 15 minutes during infusion and more frequently if signs of adverse reactions are suspected. | | | | |
| Monitor post infusion BP 30 minutes after completion of infusion. | | | | |
| Pre-Medications: | | | | |
| Acetaminophen (Tylenol) 650mg PO ONCE | | | | |
| Diphenhydramine (Benadryl) 25mg PO ONCE | | | | |
| | | | | |
| Diphenhydramine (Benadryl) 25mg IV ONCE | | | | |
| Loratadine (Claritin) 10mg PO ONCE | | | | |
| Methylprednisolone Sodium Succinate (Solu-Medrol) 125mg IV ONCE | | | | |
| Infusion Therapy: | | | | |
| Spesolimab-sbzo (Spevigo) 900mg IV over 90 minutes ONCE | | | | |
| - Flush IV line with 0.9% sodiume chloride prior to infusion | | | | |
| Administer through an IV line containing a sterile, non-pyrogenic low protein binding in-line filter (0.2 micron) | | | | |
| - Total infusion time should not exceed 180 minutes | | | | |
| - Flush IV line with 0.9% sodium chloride after infusion is complete. | | | | |
| | | | | |
| Supportive Care Medications: | | | | |
| Acetaminophen (Tylenol) 650mg PO ONCE PRN mild pain (1-3) or moderate pain (4-6). Give first if not given as a pre-medication. | | | | |
| ☑ Ibuprofen (Motrin) 800mg PO ONCE PRN mild pain (1-3) or moderate pain (4-6). Give second after acetaminophen. | | | | |
| Ondansetron (Zofran) 4mg IV ONCE PRN nausea/vomiting. | | | | |
| Hypersensitivity Protocol: | | | | |
| Initiate Atrium Health approved hypersensitivity protocol in the event of an acute adverse or anaphylactic infusion/injection reaction. The hypersensitivity protocol can | | | | |
| be found on the Atrium Health Infusion Center website at atriumhealth.org/infusion. | | | | |
| Prescriber Information: | | | | |
| Provider Name: | | Dhana | | |
| | | Phone: Fax: | | |
| Practice Name: | | NPI: | | |
| Address: | | Office Contact: | | |
| City, State, Zip: Office Contact Phone Number: | | | | |
| Physician Signature: (Order expires 12 months i | from date of signature) No St | | | |
| Signature: | | Date: | | |