

Atrium Health Infusion Center

Referral Status:	☐ New Start ☐ Order Change ☐ Renewal
Preferred Location:	Atrium Health Infusion Center Concord Fax: 704-468-3401
	Atrium Health Infusion Center Pineville Fax: 704-468-3401
	Atrium Health Infusion Center Southpark Fax: 704-468-3401
	Atrium Health Infusion Center Huntersville Fax: 704-468-3401
	Atrium Health Infusion Center Kenilworth, a facility of CMC Fax: 704-512-5390
	Atrium Health Infusion Center Abbey Place, a facility of CMC Fax: 704-512-5390
	Atrium Health Infusion Center Cabarrus, a facility of CMC Fax: 704-512-5390

Vyepti® (eptinezumab-jjmr) Infusion Order (Revised 10/16/2025)
All orders with a $\sqrt{}$ will be placed.

	·
Patient Demographics:	
Patient Name: Date of Birth:	MRN:
Address:	
City: State:	Zip Code:
Allergies: (please list all allergies or attach list)	
■ NKDA	
Diagnosis: (Complete the 2nd and/or 3rd Digits of the ICD-10)	
G43 Migraine in adults	Other:
Required Documentation: (required prior to scheduling)	
Patient Demographic Sheet	If the patient is new to the ordered therapy, indicate washout from previous therapy:
Copy of Insurance Card (front and back)	
Most Recent Labs (must include labs pertinent to medication ordered)	☐ No Washout Needed
Consult Note or last 2 Office Visits with referring provider or APP	If the patient is currently on the therapy, indicate date of last infusion: Next infusion due date:
Complete Medication List -	
Include all tried and failed meds	If this is an arder shange only indicate if the current thereny should be administered until
	If this is an order change only, indicate if the current therapy should be administered until insurance approval is received for the new request.
Diagnostic Studies Pertinent to Medication Ordered	insurance approvacts received for the new request.
Diagnostic Studies Fertillent to Medication Ordered	☐ Yes ☐ No
Nursing Communication:	
Nursing Communication: ✓ Start PIV/Access CVC and flush device per approved Atrium Health p	protocol.
✓ Start PIV/Access CVC and flush device per approved Atrium Health p✓ Obtain vital signs PRE-treatment and POST-treatment. Obtain vital s	igns PRN during treatment.
✓ Start PIV/Access CVC and flush device per approved Atrium Health p✓ Obtain vital signs PRE-treatment and POST-treatment. Obtain vital s	
✓ Start PIV/Access CVC and flush device per approved Atrium Health p✓ Obtain vital signs PRE-treatment and POST-treatment. Obtain vital s	igns PRN during treatment.
 Start PIV/Access CVC and flush device per approved Atrium Health p Obtain vital signs PRE-treatment and POST-treatment. Obtain vital s Monitor patient for signs of reaction for 30mins AFTER completion of 	igns PRN during treatment.
 ✓ Start PIV/Access CVC and flush device per approved Atrium Health process. ✓ Obtain vital signs PRE-treatment and POST-treatment. Obtain vital sreatment of Monitor patient for signs of reaction for 30mins AFTER completion of Infusion Therapy: 	igns PRN during treatment.
✓ Start PIV/Access CVC and flush device per approved Atrium Health posterior Obtain vital signs PRE-treatment and POST-treatment. Obtain vital sometime of Monitor patient for signs of reaction for 30mins AFTER completion of Infusion Therapy: Eptinezumab-jjmr (Vyepti)mg IV every 12 weeksmg IV every 12 weeksmg IV every 12 weeksmg IV every 12 weeksmg IV everymg IV everymg IV every	igns PRN during treatment. 1st infusion and subsequent infusions PRN IF previous signs of reaction observed.
✓ Start PIV/Access CVC and flush device per approved Atrium Health p ✓ Obtain vital signs PRE-treatment and POST-treatment. Obtain vital s ✓ Monitor patient for signs of reaction for 30mins AFTER completion of Infusion Therapy: ✓ Eptinezumab-jjmr (Vyepti)mg IV every 12 weeks Supportive Care Medications:	igns PRN during treatment. 1st infusion and subsequent infusions PRN IF previous signs of reaction observed.
✓ Start PIV/Access CVC and flush device per approved Atrium Health posterior Obtain vital signs PRE-treatment and POST-treatment. Obtain vital sometime of Monitor patient for signs of reaction for 30mins AFTER completion of Infusion Therapy: Eptinezumab-jjmr (Vyepti)mg IV every 12 weeksmg IV every 12 weeksmg IV every 12 weeksmg IV every 12 weeksmg IV everymg IV everymg IV every	igns PRN during treatment. 1st infusion and subsequent infusions PRN IF previous signs of reaction observed.
✓ Start PIV/Access CVC and flush device per approved Atrium Health p Obtain vital signs PRE-treatment and POST-treatment. Obtain vital s Monitor patient for signs of reaction for 30mins AFTER completion of Infusion Therapy: Eptinezumab-jjmr (Vyepti)mg IV every 12 weeks Supportive Care Medications: Ondansetron (Zofran) 4mg IV ONCE PRN for nausea/vomiting. Ibuprofen (Motrin) 800mg PO every 8 hours PRN for mild pain (1-3) of Hypersensitivity Protocol: Initiate Atrium Health approved hypersensitivity protocol in the event	igns PRN during treatment. 1st infusion and subsequent infusions PRN IF previous signs of reaction observed.
✓ Start PIV/Access CVC and flush device per approved Atrium Health process of the process of th	igns PRN during treatment. 1st infusion and subsequent infusions PRN IF previous signs of reaction observed. r moderate pain (4-6). t of an acute adverse or anaphylactic infusion/injection reaction. The hypersensitivity
✓ Start PIV/Access CVC and flush device per approved Atrium Health process of the completion of the c	igns PRN during treatment. 1st infusion and subsequent infusions PRN IF previous signs of reaction observed. r moderate pain (4-6). t of an acute adverse or anaphylactic infusion/injection reaction. The hypersensitivity
✓ Start PIV/Access CVC and flush device per approved Atrium Health postain vital signs PRE-treatment and POST-treatment. Obtain vital some Monitor patient for signs of reaction for 30mins AFTER completion of Infusion Therapy: ✓ Eptinezumab-jjmr (Vyepti)mg IV every 12 weeks Supportive Care Medications: ✓ Ondansetron (Zofran) 4mg IV ONCE PRN for nausea/vomiting. ✓ Ibuprofen (Motrin) 800mg PO every 8 hours PRN for mild pain (1-3) of Hypersensitivity Protocol: Initiate Atrium Health approved hypersensitivity protocol in the event protocol can be found on the Atrium Health Infusion Center website	igns PRN during treatment. 1st infusion and subsequent infusions PRN IF previous signs of reaction observed. r moderate pain (4-6). t of an acute adverse or anaphylactic infusion/injection reaction. The hypersensitivity
✓ Start PIV/Access CVC and flush device per approved Atrium Health p ✓ Obtain vital signs PRE-treatment and POST-treatment. Obtain vital s ✓ Monitor patient for signs of reaction for 30mins AFTER completion of Infusion Therapy: ✓ Eptinezumab-jjmr (Vyepti)mg IV every 12 weeks Supportive Care Medications: ✓ Ondansetron (Zofran) 4mg IV ONCE PRN for nausea/vomiting. ✓ Ibuprofen (Motrin) 800mg PO every 8 hours PRN for mild pain (1-3) o Hypersensitivity Protocol: ✓ Initiate Atrium Health approved hypersensitivity protocol in the even protocol can be found on the Atrium Health Infusion Center website Prescriber Information:	igns PRN during treatment. 1st infusion and subsequent infusions PRN IF previous signs of reaction observed. r moderate pain (4-6). t of an acute adverse or anaphylactic infusion/injection reaction. The hypersensitivity at atriumhealth.org/infusion.
✓ Start PIV/Access CVC and flush device per approved Atrium Health policy Obtain vital signs PRE-treatment and POST-treatment. Obtain vital signs PRE-treatment and POST-treatment. Obtain vital signs of reaction for 30mins AFTER completion of Infusion Therapy: ✓ Eptinezumab-jjmr (Vyepti)mg IV every 12 weeks Supportive Care Medications: ✓ Ondansetron (Zofran) 4mg IV ONCE PRN for nausea/vomiting. ✓ Ibuprofen (Motrin) 800mg PO every 8 hours PRN for mild pain (1-3) of Hypersensitivity Protocol: ✓ Initiate Atrium Health approved hypersensitivity protocol in the even protocol can be found on the Atrium Health Infusion Center website Prescriber Information: Provider Name:	igns PRN during treatment. 1st infusion and subsequent infusions PRN IF previous signs of reaction observed. r moderate pain (4-6). t of an acute adverse or anaphylactic infusion/injection reaction. The hypersensitivity at atriumhealth.org/infusion. Phone: Fax:
✓ Start PIV/Access CVC and flush device per approved Atrium Health postain vital signs PRE-treatment and POST-treatment. Obtain vital signs PRE-treatment and POST-treatment. Obtain vital signs of reaction for 30mins AFTER completion of Infusion Therapy: ✓ Eptinezumab-jjmr (Vyepti)mg IV every 12 weeks Supportive Care Medications: ✓ Ondansetron (Zofran) 4mg IV ONCE PRN for nausea/vomiting. ✓ Ibuprofen (Motrin) 800mg PO every 8 hours PRN for mild pain (1-3) of Hypersensitivity Protocol: ✓ Initiate Atrium Health approved hypersensitivity protocol in the even protocol can be found on the Atrium Health Infusion Center website Prescriber Information: Provider Name: Practice Name:	Igns PRN during treatment. 1st infusion and subsequent infusions PRN IF previous signs of reaction observed. r moderate pain (4-6). t of an acute adverse or anaphylactic infusion/injection reaction. The hypersensitivity at atriumhealth.org/infusion. Phone: Fax: NPI:
✓ Start PIV/Access CVC and flush device per approved Atrium Health procession of Post-treatment and Post-treatment. Obtain vital sreading Procession of Post-treatment of Summars of Post-treatment of Summars of Post-treatment. Obtain vital sreading Post-treatment of Summars of Post-treatment of Post-treatment of Summars of Post-treatment of Summars of Post-treatment of Post-t	igns PRN during treatment. 1st infusion and subsequent infusions PRN IF previous signs of reaction observed. r moderate pain (4-6). t of an acute adverse or anaphylactic infusion/injection reaction. The hypersensitivity at atriumhealth.org/infusion. Phone: Fax: NPI: Office Contact: Office Contact Phone Number: