

**Adult Renflexis Infusion Order** (Revised 4/3/21)

**Instructions to Provider:** All orders with  will be placed unless otherwise noted. Please fax completed order, along with referral form.

**Required Lab Results:** Prior to first infusion Hep B Profile and PPD/Quantiferon Gold (If outside of Atrium, fax with order. Required prior to scheduling.)

**Infusion Therapy:**

**ICD 10 code:** \_\_\_\_\_

Renflexis (infliximab-abda) \_\_\_\_\_ mg/kg IV over 2 hours (*rounded to the next 100, unless within 10% of 100mg mark then round down*)

**Frequency:** week 0, 2 and 6 then every \_\_\_\_\_ weeks (Loading)      **OR**

**Frequency:** every \_\_\_\_\_ weeks (Maintenance)

**Pre-Meds: Administer 30 minutes prior to Renflexis**

Acetaminophen 650 mg PO x 1

Benadryl \_\_\_\_\_ mg PO or \_\_\_\_\_ mg IV x 1 (*if applicable, only choose ONE*)

Loratadine 10 mg PO x 1

SoluMedrol 125 mg IV x 1

**Additional Orders:**

---

**Special Instructions:**

- **Rate for Loading Doses ( $\leq 1000$ mg dose):** 20ml/hr x 10ml, 80ml/hr x 40ml, 150ml/hr x 75ml and 250ml/hr x remainder of infusion. **Rate for maintenance dose:** 125ml/hr x 250mL.
- **Rate for Loading Doses ( $> 1000$ mg dose):** 40mL/hr x 20mL, 160mL/hr x 80mL, 300mL/hr x 150mL, 500mL/hr X remainder. **Rate for maintenance dose:** 250mL/hr x 500mL.
- Infuse using a 1.2-micron filter or less
- If patient has an infusion reaction and the Renflexis is continued per provider order, the rate will be determined by provider
- Follow Atrium Health Infusion Center protocol for hypersensitivity PRN.
- Do not administer Renflexis and notify ordering provider if patient has a temperature greater than 100°F, complains of symptoms of acute viral or bacterial illness, or if patient is taking antibiotics for current infection.
- Monitor patient for new onset or worsening congestive heart failure symptoms.

**Infusion Monitoring:**

- Obtain vital signs pre- and post-infusion. During loading doses: obtain vital signs after 1<sup>st</sup> hour of infusion and PRN.
- Monitor for signs of reaction for 30 mins after completion of 1<sup>st</sup> infusion and subsequent infusions PRN if previous signs of reaction observed

Provider Name: \_\_\_\_\_

Provider Signature: \_\_\_\_\_

Date: \_\_\_\_\_ (Order valid for 1 year)

Patient Name:

DOB:

MRN:

Atrium Health Infusion Centers  
Phone: 704-468-3400 Fax: 704-468-3401

Provider Name: \_\_\_\_\_

Provider Signature: \_\_\_\_\_

Date: \_\_\_\_\_ (Order valid for 1 year)

Patient Name:

DOB:

MRN: